

UNIVERSITY OF COPENHAGEN



Confirmation of a QTL region associated with susceptibility to ascaris suum infections in pigs

Skallerup, Per; Nejsum, Peter; Thamsborg, Stig Milan; Fredholm, Merete; Göring, Harald HH; Jørgensen, Claus Bøttcher

Publication date:
2011

Document version
Early version, also known as pre-print

Citation for published version (APA):
Skallerup, P., Nejsum, P., Thamsborg, S. M., Fredholm, M., Göring, H. HH., & Jørgensen, C. B. (2011). *Confirmation of a QTL region associated with susceptibility to ascaris suum infections in pigs*. Abstract from 23rd International Conference of the WAAVP, Buenos Aires, Argentina.

Confirmation of a QTL Region Associated With Susceptibility to *Ascaris suum* Infections in Pigs

¹Skallerup, P.; ¹Nejsum, P.; ¹Thamsborg, S. M.; ²Fredholm, M.; ³Göring, H. H. and ²Jørgensen, C. B.

¹) Parasitology, Health and Development, Department of Veterinary Disease Biology, Faculty of Life Sciences ²) Genetics & Bioinformatics, Department of Basic Animal and Veterinary Sciences, Faculty of Life Sciences, University of Copenhagen, Denmark ³) Department of Genetics, Southwest Foundation for Biomedical Research, San Antonio, TX, USA.

Abstract

Intestinal helminth infections cause health and welfare problems in both human and animal populations. Heritability analysis of a resource population of pigs infected with *Ascaris suum* (large roundworm) and *Trichuris suis* (whipworm) has shown high heritabilities for parasite burdens of both parasites. A number of putative QTL regions have been identified through a genome scan using a 7K porcine SNP-chip. To validate a QTL associated with *Ascaris* worm load on SSC4, we have genotyped two of the most promising SSC4 candidates in another population consisting of 174 pigs from 39 litters infected with *Ascaris*. The population used for the validation was produced by sows inseminated with mixed semen from three boars. The heritability (\pm SE) of worm burden of non-larval *Ascaris* (>2 cm), adult *Ascaris* (>10 cm), and total number of *Ascaris* was 73.1% (± 0.17), 31.4% (± 0.15), and 36.9% (± 0.16), respectively. The two SSC4 SNPs were genotyped using the TaqMan SNP Genotyping Assay. One SNP was associated with both adult *Ascaris* burden ($P < 0.001$) and total *Ascaris* burden ($P < 0.01$). Future research will further investigate this QTL associated with *Ascaris* burden.